

# International Journal of General Medicine & Surgery

Available Online: https://ijgms.edwiserinternational.com

# Prevalence of Acute Abdomen in General Surgery Ward of Aliabad Teaching Hospital

## Ahmadullah Danish<sup>\*</sup>

Department of abdominal surgery, Aliabad Teaching Hospital, Kabul University of Medical Sciences (Abuali Ibn Sina), Kabul, Afghanistan

#### Article info

Received 5 May 2020 Revised 7 September 2020 Published 14 September 2020

\*Corresponding author: Ahmadullah Danish, Department of abdominal surgery, Aliabad Teaching Hospital, Kabul University of Medical Sciences (Abuali Ibn Sina), Kabul, Afghanistan; E-mail: <u>dr.ahdanish@gmail.com</u>

#### Abstract

**Objective:** To determine the prevalence of acute abdomen admission and their frequency according to age and sex in general surgery ward of Aliabad teaching Hospital.

**Patients and method:** This retrospective descriptive study was performed by reviewing medical records (In-patient file) of 299 patients who admitted with diagnosis of acute abdomen between March 21, 2018 and March 21, 2019 in general surgery ward of Aliabad Teaching Hospital.

**Results:** There were 176 males (58,7%) and 123 females (41,3%), male to female ratio being 1,14:1. The mean age was 36,5 ( $\pm$ 1,09) years and the age range was 15-94 years. Highest number of patients were in 15-24 years of life (n=122, 40,8%). The most frequent cause was found to be acute appendicitis for acute abdomen accounting for 57,5% of total admission. It was followed by acute intestinal obstruction (20,7%) and acute cholecystitis (10,4%). Median age of presentation of acute appendicitis was 23 years. Median age of presentation for acute cholecystitis was 47 years. Male to female ratios in the most common cause like acute appendicitis, intestinal obstruction and acute cholecystitis were 1.5:1, 1.8:1, 0.6:1, respectively.

**Conclusion:** Acute abdomen more common in age group of 15-24 years followed by 25-34 years. Male has higher prevalence of acute abdomen when compared to female with exception of acute cholecystitis. The most common cause of acute abdomen was acute appendicitis.

Keywords: Acute abdomen; Abdominal pain; Prevalence; Acute appendicitis

## Introduction

The acute abdomen is a clinical situation in which an acute change in the condition of the intra- abdominal organs, commonly attached to inflammation or infection, needs urgent and accurate diagnosis [1]. One of the most repeatedly common symptoms in patients seeking emergency department care and the most common presenting complaint in patients with surgical diseases of the abdomen is acute abdominal pain.

According to the surgical point of view acute abdominal pain is the fundamental symptom of acute abdomen [2]. The causes of acute abdominal pain have a wide range and a detailed history, thorough clinical examination and organized investigations will guide to correct management. Evaluation of such patients and decision making for their management is very important, since some of them will have life threatening conditions that require urgent surgery which improves outcome; others may need a trial of conservative management that may require intervention if the condition fails to settle. A detailed history, full clinical examination and carefully selected investigations will guide to correct diagnosis and management [3].

Acute abdomen is a common condition encountered in emergency surgical practice [3,4]. In the developed world acute abdomen is most common in the age group 20 to 29 years with male predominance and acute abdomen accounts for 36.4% of surgical emergency [5].

The aim of this study was to identify the prevalence of acute abdomen and their frequency according to the age and sex for one year (March 21, 2018-March 21, 2019), in general surgery ward of Aliabad Teaching Hospital.

## **Method and Patients**

This retrospective descriptive study includes all adult cases that were managed as acute abdomen in the general surgery Ward of Aliabad Teaching Hospital of the Kabul University of Medical Sciences, during one year (From March 21, 2018 to March 21, 2019). Those who had traumatic injury of abdomen were excluded. Pain abdomen of urological and gynaecological origin also excluded. There were 176 males (58,7%) and 123 females (41,3%) with diagnosis of acute abdomen who underwent conservative or surgical treatment, and the age range was 15-94 years. A special data collection sheet designed to collect data regarding the diagnosis, age and sex from the profile of patients retrospectively. Data entry was performed by using Microsoft Excel 2016 and all the data were analyzed by SPSS (version 21.0). Descriptive analyses were done regarding the frequency of diseases, distribution of the diseases according to the age and sex.

## Results

The study is based on data collected from 299 patients who admitted due to acute abdomen between March 21, 2018 and March 21, 2019, including 176 males (58,7%) and 123 females (41,3%), male to female ratio being 1,14:1. Mean ( $\pm$  SE) age was 36,5 ( $\pm$  1,09) years, and the age range was 15-94 years. Highest number of patients were in 15-24 years of life (n=122, 40,8%) followed by the 25-34 years of life (n=51, 17%) (Table 1).

Total of 299 patients with acute abdomen were analyzed. The most frequent cause was found to be acute appendicitis; 57,5% of total admission, followed by acute intestinal obstruction (20,7%) and acute cholecystitis (10,4%). Other causes were acute pancreatitis (4%), peptic ulcer perforation (2%), mesenteric ischemia (1,7%), subdiaphragmatic abscess (1%), appendicular abscess (1,3%), ilium perforation (0,7%), jejunal perforation (0,3%) and peptic ulcer hemorrhage (0,3%) (Table 2).

Median age, age range and male to female ratios in the most common cause of acute abdomen like acute appendicitis, intestinal obstruction and acute cholecystitis were; (23 years, 16-78 years, 1.5:1), (50 years, 17-91 years, 1.8:1), (47 years, 16-94 years, 0.6:1), respectively (Table 3). 235 cases (78.6%) were managed surgically while 64 (21.4%) cases were managed conservatively.

Tuble 1. Age and sex distribution.						
Age group	Male, n (% of gender)	Female, n (% of gender)	Total, n (% of total)			
15-24 years	65 (36,9%)	57 (46.3%)	122 (40,8%)			
25-34 years	39 (22,2%)	12 (9,8%)	51 (17%)			
35-44 years	21 (11,9%)	25 (20,3%)	46 (15,4%)			
45-54 years	17 (9,7%)	16 (13,1%)	33 (11%)			
55-64 years	14 (8%)	8 (6,5%)	22 (7,3%)			
65-74 years	9 (5%)	2 (1,6%)	11 (3,7%)			
75-84 years	6 (3,4%)	3 (2,4%)	9 (3%)			
85-94 years	5 (2,9%)	0 (0%)	5 (1,7%)			
Total	176 (100%)	123 (100%)	299 (100%)			

Table 1: Age and sex distribution.

l'a	bl	<b>e</b> 2	2:	SĮ	pectrum	ot	disease	ın	patients	with	acute a	ιbd	lomen.	

Diagnosis	Number	Percentage (%)
Acute appendicitis	172	57.5
Acute intestinal obstruction	62	20.7
Acute cholecystitis	31	10.4

# **Citation:** Ahmadullah Danish. Prevalence of Acute Abdomen in General Surgery Ward of Aliabad Teaching Hospital. Int J Gen Med Surg 2020; 4: 130. doi: <u>10.31531/2581-8287.1000130</u>

Acute pancreatitis	12	4
Peptic ulcer perforation	6	2
Mesenteric ischemia	5	1.7
Appendicular abscess	4	1.3
Subdiaphragmatic abscess	3	1
Ilium perforation	2	0.7
Jejunal perforation	1	0.3
Peptic ulcer hemorrhage	1	0.3
Total	299	100

Table 3: Distribution of diseases according to sex and their median age with age range.

Diagnosis	Male	Female	Range (years)	Median age (years)
Acute appendicitis	103	69	16-78	23
Acute intestinal obstruction	40	22	17-91	50
Acute cholecystitis	12	19	16-94	47
Acute pancreatitis	6	6	35-60	51.5
Peptic ulcer perforation	5	1	15-80	31
Mesenteric ischemia	1	4	45-75	48
Appendicular abscess	3	1	17-50	35
Subdiaphragmatic abscess	2	1	56-85	60
Ilium perforation	2	0	23-75	49
Jejunal perforation	1	0	47	47
Peptic ulcer hemorrhage	1	0	33	33

## Discussion

In this study, frequency of acute abdomen in male was higher than female in most of the acute abdomen cases except for acute cholecystitis. In general, male to female sex ratio was 1,14:1. Male to female ratios in the most leading cause of acute abdomen like acute appendicitis, intestinal obstruction and acute cholecystitis were 1.5:1, 1.8:1 and 0.6:1, respectively. The male dominance is like what was reported in other studies [5-9]. Most patients were in their 2nd and 3rd decades of life, which was a compatibility with previous study done in other countries [6-8,10,11]. In current study we found that acute appendicitis is the most leading cause of acute abdomen accounting for 57.5% of total admission that is like other studies [7,8,11-15]. Second common cause was intestinal obstruction followed by acute cholecystitis. Similar findings were found in a study conducted by Rajesh et al. [16]. A research study was conducted by Malviya A et al where they found similar finding to our study that acute appendicitis to be most common cause of acute abdomen followed by intestinal obstruction but, they have found the hollow organ perforation more common than acute cholecystitis [17]. In contrast, our study suggests acute cholecystitis more common than perforation. Similar findings were also found other researchers in regard to frequency of acute appendicitis and intestinal obstruction but their study also showed that hollow organ perforation more common cause of acute abdomen than acute cholecystitis and this their finding also was contradicted with our study. May be, socio-economic and diet are the factors for observed difference and further researches need to find out more reasons for this difference in the future.

## Conclusion

Acute abdomen is more common in 2nd and 3rd decades of life with male predominance, exception of acute cholecystitis. The leading cause of acute abdomen in this study was acute appendicitis followed by acute intestinal obstruction and acute cholecystitis.

## Ethical approval

Ethical approval is not required for non- interventional registry-based studies in Afghanistan and no patient or public were involved.

## Funding

No funding was received for this study.

**Citation:** Ahmadullah Danish. Prevalence of Acute Abdomen in General Surgery Ward of Aliabad Teaching Hospital. Int J Gen Med Surg 2020; 4: 130. doi: <u>10.31531/2581-8287.1000130</u>

## References

- Kauffman GL. Acute abdomen in: Corson JD, Williamsons RC. Surgery. 1<sup>st</sup> Edn. Spain: Mosby International Limited 2001; 1-14.
- Vitello JM. Acute abdominal pain in: Manual of surgical therapeutics. 8<sup>th</sup> Edn. Boston: Little Brown 1993; 153.
- 3. Dang C, Aguilera P, Dang A, et al. Acute abdominal pain. Four classifications can guide assessment and management. Geriatrics 2002; 57: 30-42.
- 4. Jawaid M, Masood Z, Iqbal SA, et al. The pattern of diseases in a surgical unit at a tertiary care public hospital of Karachi. Pak J Med Sci 2004; 20: 311-314.
- 5. Tsegaye S, Osman M, Bekele A. Surgically treated acute abdomen at Gondar University Hospital, Ethiopia. East Central Afr J Surg 2007; 12: 53-57.
- Zelalem A. Pattern of acute abdomen in Yirgalem Hospital, Southern Ethiopia. Ethiop Med J 2000; 38: 227-235.
- 7. Kotiso B, Abdurrahman Z. Pattern of acute abdomen in adult's patients in Tikur Anbessa teaching hospital, Addis Ababa, Ethiopia. East Central African J Surg 2007; 12: 47-52.
- Datubo- Brown DD, Adotey JM. Pattern of surgical acute abdomen, the University of Port Harcourt teaching hospital. West Afr J Med 1990; 9: 59-62.

This manuscript was peer-reviewed Mode of Review: Single-blinded Academic Editor: Napolitano LM

- 9. Xiaoshan F, Defu L, Shuchang C. Analysis of the causes of acute surgical abdomen in Nazareth, Ethiopian. Annual medical conference of the Ethiopian Medical Association (EMA), 1998.
- 10. Adesunkanmi AR. Changing pattern of acute intestinal obstruction in a Tropical African Population. East Afr Med J1996; 73: 727-731.
- 11. Ogbonna BC, Obekpa PO, Momoh JT, et al. Another look at acute appendicitis in tropical Africa and the value of laparoscopy in diagnosis. Trop Doc 1993; 23: 82-84.
- 12. Ayalew T. Small intestinal volvulus in adults of Gondar region. Ethiopia Med J 1992; 30: 111-114.
- 13. Out AA. Tropical surgical abdominal emergencies: Acute appendicitis. Trop Geogr Med 1989; 41: 188-122.
- 14. Ajao OG. Abdominal emergencies in a tropical African population. Br J Surg 1981; 68: 345-347.
- 15. Okoro IO. Appendicitis in Zaire: A study of surgical epidemiology, etiology and bacteriology and the leukocyte pattern in appendicitis. National Postgraduate Medical College of Nigeria FMCS part II Dissertation 1987
- Poudel E. Prevalence of acute abdomen in surgery ward at Nepal. J Universal College Med Sci 2019; 19: 14-15.
- 17. Malviya A, Hussain A, Bulchandani HP, et al. A comprehensive study on acute non-traumatic abdominal emergencies. Int Surg J 2017; 22: 4: 2297-2302.

**Copyright:** ©2020 Ahmadullah Danish. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<u>http://creativecommons.org/licenses/by/4.0/</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.